

How black holes turn cusps into cores

Marc Hemsendorf (marchems@physics.rutgers.edu)

*Dept. of Physics and Astronomy, Rutgers University, 136 Frelinghuysen Rd.,
Piscataway, NJ 08854, USA*

Abstract. Collapsing collisionless particle systems form gravitational bound halos with cuspy density profiles. Also hierarchical merging of these systems produce remnants with cuspy central density profiles. These results lead to the assumption of cuspy NFW [5] profiles for the density distribution in dark matter halos. However, observed rotation curves in disk galaxies suggest dark matter halos with isothermal core. The same kind of problem can be found for globular clusters, which show cores in their density profiles but should be cuspy if they were formed through a cold collapse.[4, 3] We are showing how small, massive, and compact objects can efficiently transform cuspy stellar systems into density profiles with an isothermal core.

Keywords: black holes, globular clusters, rotation curves, dark matter halos

1. Introduction

2. Numerical experiments

2.1. INITIAL CONDITIONS

For the realizations of cuspy systems we are choosing Dehnen models with isotropic velocity dispersion. The Dehnen family of models are characterized by a parameter γ that measures the degree of central concentration.[2] The density profile is

$$\rho(r) = \frac{(3 - \gamma)M}{4\pi} \frac{a}{r^\gamma (r + a)^{4-\gamma}}, \quad (1)$$

M being the total mass and a being the scale length. Both M and a are set to unity. We chose centrally condensed Dehnen models with inner slope $\gamma = 1, 3/2$, and 2 , yielding a density profile similar to those of elliptical galaxies with dense stellar nuclei. The black holes and the stars are initially distributed according to the same profile.

We are creating several realizations of these Dehnen models with a total particle number of 9910. These realizations contain 10 black holes with a combined mass of 0.01 which is 1% of the total mass of the system.



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.




Figure 1. Comparison of the initial radial density distribution of the stars with the average radial density distribution of the stars after 600 N -body time units.

2.2. METHOD

We are using NBODY6++ for the orbit integration. NBODY6++ is a high precision fourth order direct force integrator [6, 1]. Since it can regularize the equations of motion for particles having close encounters no softening needs to be applied. This is why we are able to follow the formation of black hole – black hole binaries transferring part of their binding energy into the stellar system.

We stopped the simulations once the number of black holes bound to the stellar system dropped below three. This process usually took a few hundred dynamical times.

2.3. RESULTS

Figures 1 and 2 summarize our results. Both plots compare the initial model with the situation after 600 N -body time units. This corresponds to an evolution over roughly 100 dynamical timescales. We have averaged the results of four realizations in order to find $\rho(r)$ and $\sigma(r)$ in Figures 1 and 2.

Figure 1 shows the initial radial density distribution compared with the averaged radial density distribution after 600 N -body time units. While the impact of the black holes removes mass up to radius r_{crit}




Figure 2. Comparison of the initial 3D velocity dispersion of the stars with the average 3D velocity dispersion of the stars after 600 N -body time units.

Table I. The amount of relocated mass after 600 N -body time units

γ	r_{crit}	M_{rel}
1	0.5	0.06
3/2	0.4	0.07
2	0.3	0.07

of 0.5 which is half way to the initial scale radius for $\gamma = 1.0$ and $\gamma = 1.5$, the effect is seemingly smaller in the models with $\gamma = 2.0$. As Table I shows, the amount of mass relocated by the black holes remains constant.

Figure 2 shows the initial 3D velocity dispersion in the models and the 3D velocity dispersion after 600 N -body time units. Since models with $\gamma = 2.0$ are isothermal, the evolution is quite modest for such models. In the case of smaller γ , Dehnen models have a zero velocity dispersion in the center. Our models with smaller γ show a strong evolution into isothermal systems through the influence of the black holes.

3. Discussion

We have shown that 10 black holes are capable of efficiently destroying a stellar cusp within a few ten dynamical timescales. In our simulations each black hole removes an equivalent of seven times its own mass from the central regions of the system. The resulting systems show isothermal cores.

Acknowledgements

This work was supported by NSF grant 00-71099 and by NASA grants NAG5-6037 and NAG5-9046. I would like to thank D. Merritt and R. Jimenez for their encouragement and help. I am grateful to the NASA Center for Computational Sciences at NASA-Goddard Space Flight Center, the National Computational Science Alliance, the Center for Advanced Information Processing at Rutgers University, the John von Neumann Institut für Computing in Jülich, and the Höchstleistungsrechenzentrum in Stuttgart for their generous allocations of computer time.

References

1. Aarseth, S. J.: 1999, ‘From NBODY1 to NBODY6: The growth of an industry’. *Publications of the astronomical society of the pacific* **111**, 1333–1346.
2. Dehnen, W.: 1993, ‘A family of potential–density pairs for spherical galaxies and bulges’. *Monthly Notes of the Royal Astronomical Society* **265**, 250–256.
3. Jenkins, A., C. S. Frenk, F. R. Pearce, P. A. Thomas, J. M. Colberg, S. D. M. White, H. M. P. Couchman, J. A. Peacock, G. Efstathiou, and A. H. Nelson: 1998, ‘Evolution of structure in cold dark matter universes’. *The Astrophysical Journal* **499**, 20–40.
4. King, I.: 1962, ‘The structure of star clusters. I. An empirical density law.’. *The Astronomical Journal* **67**(8), 471–485.
5. Navarro, J. F., C. S. Frenk, and S. D. M. White: 1996, ‘The structure of cold dark matter halos’. *The Astrophysical Journal* **462**, 563–575.
6. Spurzem, R. and H. Baumgardt: 1999, ‘A parallel implementation of an Aarseth *N*-body integrator on general and special purpose supercomputers’. ARI-Preprint No. 68, submitted to Monthly Notes of the Royal Astronomical Society.